

Anomalies of mesosphere/lower thermosphere parameters during solar minimum 23/24

Ch. Jacobi¹, P. Hoffmann¹, M. Placke², G. Stober²

¹Institute for Meteorology, University of Leipzig, Stephanstr. 3, 04103 Leipzig, Germany

²Leibniz Institute of Atmospheric Physics at the Rostock University, Schloßstraße 6, 18225 Kühlungsborn, Germany

Abstract

The recent solar minimum has been characterized by an anomalous strong decrease in thermospheric density since 2005. Here we analyze anomalies of mesosphere/lower thermosphere parameters possibly connected with this effect. In particular, nighttime mean LF reflection heights measured at Collm, Germany, show a very strong decrease after 2005, indicating a density decrease. This decrease is also visible in mean meteor heights measured with a VHF meteor radar at Collm. This density decrease is accompanied by an increase of gravity wave (GW) amplitudes in the upper mesosphere and a decrease in the lower thermosphere. On the decadal scale, GW are negatively correlated with the background zonal wind, but this correlation is modulated in the course of the solar cycle, indicating the combined effect of GW filtering and density decrease.

1 Introduction

It is widely known that solar variability influences the atmosphere (Gray et al., 2010), e.g. the dynamics of the middle and upper atmosphere. In particular, search for an effect of the 11-year solar Schwabe cycle has been undertaken, for example, to explain part of the observed variability of mesosphere and lower thermosphere (MLT), which can be studied by radars. Indeed, indication for a solar effect has been found in MLT radar wind time series over Central Europe (Jacobi and Kürschner, 2006; Keuer et al., 2007).

Solar cycles are different from one to another. Especially, the recent solar minimum has been extremely extended and extraordinarily deep. Consequently, it led to extreme upper atmosphere reactions, in particular a decrease in thermospheric density (Emmert et al., 2010; Solomon et al., 2010) which exceeds the expectations that would have been based on conventional solar indices like the sunspot number or F10.7.

Lower ionospheric electron density reacts on the solar cycle, which leads to an 11-year modulation of radio wave reflection heights. These have been observed, e.g., by Kürschner and Jacobi (2003) who found that the Collm LF reflection heights (177 kHz, distance to transmitter about 160 km) are about 2 km lower during solar maximum than during solar minimum. This is owing to the increased ionisation during solar maximum, however, this effect is superposed by thermal shrinking of the mesosphere during solar minimum, since the middle atmosphere has a solar cycle

signal of about 2 K difference between maximum and minimum. However, this thermal shrinking is usually overcompensated by increased ionisation. We are interested, whether the recent extreme solar minimum has led to anomalous signatures either in MLT wind or density. We focus on the summer MLT, which is not that much influenced by stratospheric planetary wave activity.

2 Measurements

2.1 Collm LF lower ionospheric drifts, reflection heights, and GW estimates

At Collm Observatory, MLT winds have been obtained by D1 LF radio wind measurements from 1959-2008, using the ionospherically reflected sky wave of three commercial radio transmitters. The data are combined to half-hourly zonal and meridional mean wind values. The virtual reflection heights have been estimated since late 1982 using measured travel time differences between the separately received ionospherically reflected sky wave and the ground wave (Kürschner et al., 1987). More details of the Collm LF system are given in Jacobi (2010).

Since the LF reference height changes in the course of the day, a continuous time series is not available and consequently GW spectra cannot be calculated. However, using the method presented by Gavrilov et al. (2001), horizontal wind fluctuations in the period range of 0.7-3 h can be obtained which may serve as GW wave proxy. Jacobi et al. (2006) has used this method to analyze the Collm dataset from 1984–2003. They found an 11-year solar cycle with larger GW amplitudes during solar maximum, but their dataset did not include the recent solar minimum.

2.2 Collm meteor radar

At Collm Observatory (51.3°N, 13°E), a SKiYMET meteor radar (MR) is operated on 36.2 MHz to measure horizontal winds, meteor rates and heights, and further meteor parameters since August 2004. The radar and the hourly wind detection is described in Jacobi (2010). Monthly mean wind parameters are obtained from half-hourly mean winds applying a multiple regression analysis including the mean wind, and tidal components. Based upon 2-hourly means, GW variances and fluxes are obtained by fitting the 2-hourly mean GW fluxes to the radial drift variances according to Hocking (2005). Details can be found in Placke et al. (2010).

2.3 GW potential energy from SABER

The SABER instrument on the TIMED satellites (Russell et al., 1999; Mertens et al., 2001; 2004) scans the atmosphere from about 52° of one hemisphere to 83° of the other. This latitude range is reversed by a yaw manoeuvre every 60-days. Due to the sun-synchronous orbital geometry the spacecraft passes the equator always at the same local time (12LT) on the day side. Each single temperature profile, having a vertical resolution of 0.5 km, is high-pass filtered to analyze waves with vertical wavelength of up to 6 km. From these filtered data, the vertical structure of GW amplitudes and their specific potential energy is obtained. This method has frequently been applied to GPS radio occultations (e.g., Fröhlich et al. 2007). The total energy

integrated over a sliding vertical column (10 km) is used to study GW. Note that limb scanning of the atmosphere by SABER only reveals certain parts of the GW spectrum due to the integration along the line of sight (Preusse et al., 2006). Another limitation is made by the chosen vertical filter, which allows only GW of short wavelengths to be studied.

3 GW proxy decadal variability

Jacobi et al. (2006) had shown that there is a solar cycle influence on GW activity as measured by LF over Collm. Figure 1 presents the summer (JJA) mean time series of GW variance $\zeta'^2 = u'^2 + v'^2$ at 100 km virtual height, which represents approximately 91 km real height (Jacobi, 2010). The data are an update from Jacobi et al. (2006). The 13-monthly mean sunspot number is added. Clearly, there is a solar cycle in the GW variance, and from visual inspection a decreasing long-term trend is also visible. We thus added, as a red solid line, a least squares fit of a linear trend superposed by a solar cycle

$$\zeta'^2 = a + b \times \text{yr} + c \times R \quad , \quad (1)$$

with R being the 13-monthly smoothed relative sunspot number. We are only interested in long-term and qualitative connections, so employing other widely used solar proxies as F10.7 are not superior to use of R here. As can be seen in the lower part of the figure, the residuals are not normally distributed, and generally the model is not valid during the 1996 as well as the recent solar minimum. Using the same model, but including only data until 1995 (blue line) reveals that the time interval until the early 2000s is well represented by that model, and after 2004 there is a drastic change. We therefore conclude that there is a change in dynamical regime since about 2004.




Figure 1: JJA mean LF GW proxy, and fit including linear trend and a solar cycle according to Eq. (1) at 100 km virtual height (approx. 91 km real height). The fit was performed both using the complete dataset 1984-2007 (red curve) and using part of the dataset until 1995 (blue curve). In the lower part the respective residuals are given as dashed lines. The sunspot number is also added as green line.




Figure 2: JJA mean LF GW proxy (black) and zonal mean wind (blue) at 100 km virtual height. In the lower part of the figure, running correlation coefficients between GW proxy and mean wind are added. The sunspot number is added as green line.

In the case of saturated GW, linear theory predicts amplitudes proportional to the intrinsic phase speed. Consequently, since GW phase speeds must be positive (eastward) in the summer MLT owing to the filtering effect of the stratospheric and mesospheric easterlies, a negative correlation is expected between the background wind and the GW amplitudes. In Figure 2 we present GW variance together with the background mean zonal wind at 100 km virtual height. Note that the background wind is simply the mean of the zonal wind averaged over those times when GW amplitudes have been calculated, and thus may deviate from the prevailing wind. There is an overall anticorrelation between GW variances and zonal winds, as expected from linear theory. However, during solar minimum the correlation reverses. The running correlation (Kodera, 1993) between GW proxy and mean zonal wind is added in the lower part of Figure 2. Due to the shortness of the time series, only 5 data points are used for each calculation. A clear solar cycle modulation is visible. The running correlation is correlated with the sunspot number time series with a correlation coefficient of $r = -0.80$. It is also remarkable that this modulation takes place during each solar minimum since 1986. Note also that the increase of GW variances in 2005, when solar flux already decreases and decreasing GW variances are expected, has its counterpart in a peak in 1993. We may conclude that there is obviously a different regime of mean wind-GW coupling during solar minimum, which is, however, more emphasized during the recent minimum.

LF height measurements at Collm have stopped after 2007, so that the solar minimum is not completely covered by them. To analyze winds and waves during the minimum, in Figure 3 MR summer mean zonal prevailing winds at 6 height gates are presented. Clearly, interannual variability of winds in the upper and lower height gates is opposite. This is explained by GW acceleration and filtering in the mesosphere. In the case of strong/weak mesospheric easterlies, GW amplitudes are large/small, which then lead to strong/weak vertical wind shear. Figure 3 shows that above 91 km winds

are decreasing until 2007, which is qualitatively consistent with the decrease of the LF winds during 2005-2007. After 2007, winds are increasing again. This agrees with the zonal LF wind decrease after 1994 (Figure 2). This LF zonal wind decrease is accompanied by GW proxy amplitude decrease. MR GW analyses of the recent minimum (Figure 4) again show this effect qualitatively. The same is the case for SABER potential energy (Figure 5), although here only the GW amplitudes above ~ 100 km decrease with time. This may be due to the fact that we present zonal mean potential energy, while Figures 1-4 represent point measurements and non-zonal structures are likely to exist in the MLT. Note that the above mentioned trends are only valid for heights above 90 km, while for the lower height gates the winds and GW amplitudes behave in an opposite manner.




Figure 3: Collm June-August mean zonal prevailing winds measured by meteor radar.




Figure 4: Horizontal wind variance calculated within 2-hr intervals using Collm MR wind measurements for 4 height gates.




Figure 5: JJA mean potential energy at 45°N from SABER temperature profiles. Data are averages over a 10 km vertical window, and means over all longitudes.

5 LF virtual heights and mean meteor altitudes

The solar modulation of correlation between zonal wind and GW amplitudes suggests that the decrease of reference height of wind systems may play a role. Collm LF virtual nighttime (22-2LT) reflection heights are shown in Figure 6. A multiple linear fit after Eq. (1), but analyzing virtual height instead of variance is added as red line, as well as the residuals (red dashed line). The sunspot number is added as green line. Note that real heights and height differences are much smaller than virtual height differences. Thus, the strong decrease of LF heights after 2005, for example, represents a real height decrease of 2 km only. From the residuals in Figure 6 one can see that the recent minimum is outstanding. Generally, there is an ionization driven solar cycle of reflection heights such that these are lower during solar maximum than during solar minimum. Thus, the strong decrease after 2005 is unexpected. However, a similar variability has already been observed after 1993 during the last solar minimum, although with much smaller amplitudes, LF reflection height variability is influenced by changes in ionization and mesospheric shrinking. According to Figure 6 this would mean that during the recent solar minimum thermal shrinking has overcompensated the ionization effect.

Note that the reference height changes shown in Figure 6 do not show the real variations of a line of constant pressure, for example. A better proxy is the mean height of meteors because these, constant meteor parameters as mass and velocity assumed, burn at a height that is determined by the density distribution. Figure 7 shows June-August mean meteor heights from 2005-2010, plotted against the solar flux. As expected, the meteor heights increase with solar activity. Note, however, that the mean meteor heights during 2008 and 2010, at the same level of solar flux, are different. This is partly due to the fact that the decrease is superposed by the long-term changes of mesospheric density. Bremer and Peters (2008) found a long-term decrease of -30 m/yr from LF reflection heights (and excluding the solar cycle). Subtracting this

from the measured meteor heights in Figure 7 (red line) shows that then the 2008 and 2010 heights, at the same solar flux, have exactly the same values when long-term cooling of the middle atmosphere is taken into account.

There remains some sort of hysteresis, so that the density increases about one year later that the solar flux does. A delay of one year of noctilucent cloud occurrence and solar activity has been reported by DeLand et al. (2006) and Bremer et al. (2009). Ortiz de Adler and Elias (2008) showed a similar hysteresis in ionospheric foF2 data. Jacobi et al. (2008) showed that MLT planetary wave activity lags the solar cycle by 1-3 years.




Figure 6: Collm LF virtual nighttime (22-2LT) reflection heights. A linear fit according to Eq. (1) is added as red line, as well as the residuals (red dashed line). The sunspot number is added as green line.




Figure 7: JJA mean meteor heights over Collm vs. F10.7 solar flux. The red curve represents meteor heights after removing a linear trend of -30 myr^{-1} .

6 Discussion and conclusions

Linear theory predicts that GW amplitudes are proportional to the intrinsic phase speed and thus, in the case of a given GW with specified phase speed, by the zonal wind itself. The positive correlation between GW amplitudes and zonal winds during solar minimum is thus unexpected at first glance, but may be explained by a downward shift of the GW maximum owing to thermal shrinking of the MLT wind systems. In such a case, GW already maximize in the upper mesosphere. Then, strong/weak mesospheric easterlies, which are connected with weak/strong lower thermospheric westerlies, are connected with large/small GW amplitudes, large/small GW drag and consequently strong/weak westerly winds at greater altitudes. This may explain the positive correlation during solar minimum, while there is a negative correlation during the other years, then simply in accordance with linear theory.

Whether or not the above mentioned coupling processes really work requires more detailed analyses, including more satellite analyses, further radars, and numerical modeling. However, the observations of MLT GW, mean winds, and reference heights already suggest that there is a height shift during solar minimum which may influence vertical coupling between mesosphere and lower thermosphere. The recent solar minimum represents an extreme case, but the fundamental variability, as shown by the LF measurements, was not qualitatively (although quantitatively) different from the last solar minimum, at least as far as the LF measurements at Collm are concerned. In the thermosphere, however, density decrease during the recent minimum was extreme. Thus, there are still open questions concerning solar variability and its effect on the MLT.

Acknowledgements

This study was supported by Deutsche Forschungsgemeinschaft under JA 836/22-1.

References

- Bremer, J., and Peters, D., 2008: Influence of stratospheric ozone changes on long-term trends in the meso- and lower thermosphere. *J. Atmos. Solar.-Terr. Phys.*, 70, 1473-1481.
- Bremer, J., Hoffmann, P., Latteck, R., Singer, W., and Zecha, M., 2009: Long-term changes of (polar) mesosphere summer echoes. *J. Atmos. Solar.-Terr. Phys.*, 71, 1571-1576.
- DeLand, M.T., Shettle, E.P., Thomas, G.E., and Olivero, J. J., 2006: A quarter-century of satellite polar mesospheric cloud observations. *J. Atmos. Solar.-Terr. Phys.*, 68, 9-29.
- Emmert, J.T., Lean, J.L., and Picone, J.M., 2010: Record-low thermospheric density during the 2008 solar minimum. *Geophys. Res. Lett.*, 37, L12102, doi:10.1029/2010GL043671.
- Fröhlich, K., Schmidt, T., Ern, E., Preusse, P., de la Torre, A., Wickert, W., and Jacobi, Ch., 2007: The global distribution of gravity wave energy in the lower stratosphere derived from GPS data and gravity wave modelling: Attempt and challenges., *J. Atmos. Solar.-Terr. Phys.*, 69, 2238-2248 (2007).

- Gavrilov, N.M., Jacobi, Ch., and Kürschner, D., 2001: Short-period variations of ionospheric drifts at Collm and their connection with the dynamics of the lower and middle atmosphere. *Phys. Chem. Earth*, 26, 459-464.
- Gray, L. J., Beer, J., Geller, M., Haigh, J. D., Lockwood, M., Matthes, K., Cubasch, U., Fleitmann, D., Harrison, G., Hood, L., Luterbacher, J., Meehl, G.A., Shindell, D., van Geel, B., and White, W., 2010: Solar influences on climate. *Rev. Geophys.*, 48, RG4001, doi:10.1029/2009RG000282.
- Hocking, W.K., 2005: A new approach to momentum flux determinations using SKiYMET meteor radars. *Ann. Geophys.* 23, 2433–2439.
- Jacobi, Ch., and Kürschner, D., 2006: Long-term trends of MLT region winds over Central Europe. *Phys. Chem. Earth*, 31, 16-21.
- Jacobi, Ch., Gavrilov, N.M., Kürschner, D., and Fröhlich, K., 2006: Gravity wave climatology and trends in the mesosphere/lower thermosphere region deduced from low-frequency drift measurements 1984-2003 (52.1°N, 13.2°E). *J. Atmos. Solar-Terr. Phys.*, 68, 1913-1923.
- Jacobi, Ch., Hoffmann, P., and Kürschner, D., 2008: Trends in MLT region winds and planetary waves, Collm (52°N, 15°E). *Ann. Geophys.*, 26, 1221-1232.
- Jacobi, Ch., 2010: Meteor radar measurements of mean winds and tides over Collm (51.3°N, 13°E) - comparison with LF drift measurements 2005-2007. Submitted to *Adv. Radio Sci.*
- Keuer, D., Hoffmann, P., Singer, W., and Bremer, J., 2007: Long-term variations of the mesospheric wind field at mid-latitudes. *Ann. Geophys.*, 25, 1779–1790.
- Kodera, K., 1993: Quasi-decadal modulation of the influence of the equatorial quasi-biennial oscillation on the north polar stratospheric temperatures. *J. Geophys. Res.*, 98, 7245-7250.
- Kürschner, D., Schminder, R., Singer, W., and Bremer, J., 1987: Ein neues Verfahren zur Realisierung absoluter Reflexionshöhenmessungen an Raumwellen amplitudenmodulierter Rundfunksender bei Schrägeinfall im Langwellenbereich als Hilfsmittel zur Ableitung von Windprofilen in der oberen Mesopausenregion. *Z. Meteorol.*, 37, 322-332.
- Kürschner, D., and Jacobi, Ch., 2003: Quasi-biennial and decadal variability obtained from long-term measurements of nighttime radio wave reflection heights over central Europe. *Adv. Space Res.* 32, 1701-1706, doi: 10.1016/S0273-1177(03)00773-2.
- Mertens, C. J., Mlynczak, M. G., Lopez-Puertas, P. P., Wintersteiner, R. H., Picard, J. R., Winick, L. L., Gordley, and J. M. Russell III, 2001: Retrieval of mesospheric and lower thermospheric kinetic temperature from measurements of CO₂ 15 μm earth limb emission under non-LTE conditions. *Geophys. Res. Lett.*, 28, 1391–1394.
- Mertens, C. J., Schmidlin, F.J., Goldberg, R.A., Remsberg, E.E., Pesnell, W.D., Russell, J.M. III, Mlynczak, M.G., López-Puertas, M., Wintersteiner, P.P., Picard, R.H., Winick, J.R., Gordley, L.L., 2004: SABER observations of mesospheric temperatures and comparisons with falling sphere measurements taken during the 2002 summer MaCWAVE campaign. *Geophys. Res. Lett.*, 31, L03105, doi:10.1029/2003GL018605.

- Ortiz de Adler, N. , and Elias, A.G., 2008: Latitudinal variation of foF2 hysteresis of solar cycles 20, 21 and 22 and its application to the analysis of long-term trends. *Ann. Geophys.*, 26, 1269–1273.
- Placke, M., Stober, G., and Jacobi, Ch., 2010: Gravity wave momentum fluxes in the MLT—Part I: Seasonal variation at Collm (51.3°N, 13.0°E). *J. Atmos. Solar–Terr. Phys.*, in press, doi:10.1016/j.jastp.2010.07.012.
- Preusse, P., Ern, M., Eckermann, S.D., Warner, C.D., Picard, R.H., Knieling, P., Krebsbach, M., Russell, J.M., Mlynczak, M.G., Mertens, C.J., and Riese, M., 2006: Tropopause to mesopause gravity waves in August: measurement and modelling, *J. Atmos. Solar.-Terr. Phys.*, 68, 1730–1751.
- Russell, J.M. III, Mlynczak, M.G., Gordley, L.L., Tansock, J., and Esplin, R.: 1999: An overview of the SABER experiment and preliminary calibration results. In *Proceedings of the SPIE*, 3756, 44th Annual Meeting, Denver, Colorado, July 18–23, 277–288.
- Solomon, S.C. , Woods, T.N., Didkovsky, L.V., Emmert, J.T., and Qian, L., 2010: Anomalously low solar extreme-ultraviolet irradiance and thermospheric density during solar minimum. *Geophys. Res. Lett.*, 37, L16103, doi:10.1029/2010GL044468.