

---

# Quantum Field Theory of Many-Particle Systems - Problem Set

## 10

---

*Summer Term 2014*

**Internet:** You can download this problem set at <http://www.uni-leipzig.de/~rosenow>.

### 14. Josephson effect

*4+4+4+4+4 Punkte*

Electron tunneling between two superconductors has two kinds of contributions. One is due to tunneling of single particles, as in the case between two metals or between metal and superconductor. The other is tunneling of pairs, which gives rise to the Josephson effect. The notation is the same as in problem 13. There, we showed that the single particle tunnel current can be expressed as

$$I_{\text{single}} = e \int_{-\infty}^{\infty} dt' \Theta(t-t') \left\{ e^{ieV(t'-t)} \langle [A(t), A^\dagger(t')] \rangle - e^{ieV(t-t')} \langle [A^\dagger(t), A(t')] \rangle \right\} .$$

where the operator  $A$  is defined as

$$A = \sum_{\mathbf{k}, \mathbf{p}} T_{\mathbf{k}, \mathbf{p}} c_{\mathbf{k}}^\dagger c_{\mathbf{p}} .$$

In the case of tunneling between two superconductors, there is another contribution

$$I_{\text{pair}}(t) = e \int_{-\infty}^{\infty} dt' \Theta(t-t') \left\{ e^{-ieV(t+t')} \langle [A(t), A(t')] \rangle - e^{ieV(t+t')} \langle [A^\dagger(t), A^\dagger(t')] \rangle \right\} .$$

In the following, we evaluate the voltage and time dependence of this pair contribution to the tunnel current.

- a) The most unusual feature of this expression is the fact that the time dependence in the exponential functions is  $t+t'$ . Rewrite this as  $2t+t'-t$  and change the variable of integration to  $t'' = t - t'$ . Show that the pair current can be expressed as

$$I_{\text{pair}} = 2e \operatorname{Im} \left[ e^{-2eiVt} C_{A,A}^+(eV) \right] .$$

Here,  $C_{A,A}^+(eV)$  is the Fourier transform of the retarded correlation function

$$C_{A,A}^+(t) = -i\Theta(t) \langle [A(t), A(0)] \rangle .$$

- b) In order to calculate the retarded correlation function  $C_{A,A}^+(eV)$ , we start from the Matsubara function

$$C_{A,A}^\tau(i\omega_n) = \int_0^\beta d\tau e^{i\omega_n \tau} \langle T_\tau A(\tau) A(0) \rangle .$$

Show that

$$C_{A,A}^\tau(i\omega_n) = 2 \sum_{\mathbf{k},\mathbf{p}} T_{\mathbf{k},\mathbf{p}} T_{-\mathbf{k},-\mathbf{p}} T \sum_{\epsilon_l} F_L^\dagger(i\epsilon_l, \xi_{\mathbf{k}}) F_R(i\epsilon_l - i\omega_n, \xi_{\mathbf{p}}) .$$

Here,

$$F(i\epsilon_l, \xi_{\mathbf{p}}) = \langle c_{-\mathbf{p},\downarrow}(-i\epsilon_l) c_{\mathbf{p},\uparrow}(i\epsilon_l) \rangle , \quad F^\dagger(i\epsilon_l, \xi_{\mathbf{p}}) = \langle c_{\mathbf{p},\uparrow}^\dagger(-i\epsilon_l) c_{-\mathbf{p},\downarrow}^\dagger(i\epsilon_l) \rangle$$

are the (1,2) and the (2,1) element of the Gorkov Green function matrix

$$\mathcal{G}(i\epsilon_l, \xi_{\mathbf{p}}) = \frac{1}{(i\epsilon_l)^2 - \xi_{\mathbf{p}}^2 - |\Delta_0|^2} \begin{pmatrix} i\epsilon_l + \xi_{\mathbf{p}} & -\Delta_0 \\ -\bar{\Delta}_0 & i\epsilon_l - \xi_{\mathbf{p}} \end{pmatrix} .$$

Note that for the evaluation of  $F_L^\dagger(i\epsilon_l, \xi_{\mathbf{k}})$  the order parameter in the Gorkov Green function is  $\bar{\Delta}_L$  instead of  $\bar{\Delta}_0$ , and  $\Delta_R$  instead of  $\Delta_0$  for the evaluation of  $F_R(i\epsilon_l, \xi_{\mathbf{p}})$ .

c) Use spectral representations

$$F^\dagger(i\epsilon_l, \xi_{\mathbf{k}}) = \bar{\Delta}_L \int \frac{d\omega}{2\pi} \frac{A(\omega, \xi_{\mathbf{k}})}{i\epsilon_l - \omega}$$

and

$$F(i\epsilon_l, \xi_{\mathbf{p}}) = \Delta_R \int \frac{d\omega}{2\pi} \frac{A(\omega, \xi_{\mathbf{p}})}{i\epsilon_l - \omega}$$

for both Green functions to evaluate the Matsubara sum in the expression for  $C_{A,A}^\tau(i\omega_n)$ . You will have to perform a partial fraction decomposition and to make use of the identity

$$T \sum_{\epsilon_l} \frac{e^{i\eta\epsilon_l}}{i\epsilon_l - \xi_{\mathbf{k}}} = n_F(\xi_{\mathbf{k}}) .$$

Show that  $C_{A,A}^\tau(i\omega_n)$  is given by

$$C_{A,A}^+(i\omega_n) = 2\bar{\Delta}_L \Delta_R \sum_{\mathbf{k},\mathbf{p}} T_{\mathbf{k},\mathbf{p}} T_{-\mathbf{k},-\mathbf{p}} \int_{-\infty}^{\infty} \frac{d\epsilon}{2\pi} \frac{d\epsilon'}{2\pi} A^*(\epsilon, \xi_{\mathbf{k}}) A(\epsilon', \xi_{\mathbf{p}}) \frac{n_F(\epsilon) - n_F(\epsilon')}{i\omega_n + \epsilon - \epsilon'} .$$

d) Show that the spectral function depends on  $\xi_{\mathbf{k}}$  only via  $\lambda_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_0|^2}$ , and that it is given by

$$A(\epsilon, \lambda_{\mathbf{k}}) = 2\pi \frac{1}{2\lambda_{\mathbf{k}}} [\delta(\epsilon - \lambda_{\mathbf{k}}) - \delta(\epsilon + \lambda_{\mathbf{k}})] .$$

Use this expression for the spectral function and make use of the momentum independence of tunnel matrix elements. Show that in the limit of zero temperature and after analytical continuation  $i\omega_n \rightarrow \omega + i\eta$  the retarded correlation function is given by

$$C_{A,A}^+(eV) = \frac{1}{2} |\bar{\Delta}_L \Delta_R T_0^2| e^{i\varphi} \sum_{\mathbf{k},\mathbf{p}} \frac{1}{\lambda_{\mathbf{k}} \lambda_{\mathbf{p}}} \left[ \frac{1}{eV - \lambda_{\mathbf{k}} - \lambda_{\mathbf{p}}} - \frac{1}{eV + \lambda_{\mathbf{k}} + \lambda_{\mathbf{p}}} \right] .$$

- e) Assume now that  $|\Delta_L| = |\Delta_R| \equiv \Delta_0$ . Convince yourself that for  $eV < 2\Delta_0$  the  $\delta$ -function part in  $C_{A,A}^+$  vanishes and that for the same reason there are no singular contributions to the momentum sums. As a consequence, one can write

$$C_{A,A}^+(eV) = \frac{1}{2e} J_S(eV) e^{i\varphi} ,$$

where  $J_S(eV)$  depends smoothly on voltage for  $eV \rightarrow 0$ . Show that in this notation the pair contribution to the tunneling current is given by

$$I_{\text{pair}} = J_S(eV) \sin(\omega t + \varphi) \quad \text{with} \quad \omega = \frac{2eV}{\hbar} .$$