

---

## Advanced Quantum Mechanics - Problem Set 7

---

*Winter Term 2019/20*

**Due Date:** Hand in solutions to problems marked with \* before the lecture on **Friday, 06.12.2019, 09:15**. The problem set will be discussed in the tutorials on Wednesday, 11.12.2019, and Friday, 13.12.2019.

### \*17. Continuity equation for the Dirac equation

*5 Points*

Prove the continuity equation

$$\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{j} = 0,$$

with

$$\mathbf{j} = \Psi^\dagger \begin{pmatrix} 0 & \boldsymbol{\sigma} \\ \boldsymbol{\sigma} & 0 \end{pmatrix} \Psi,$$

and  $\rho = \Psi^\dagger \Psi$  for all solutions  $\Psi$  of the Dirac equation.

### \*18. Free particle solutions of the Dirac equation

*3 Points*

Calculate the eigenvalues of the free-particle Dirac equation

$$\begin{pmatrix} m & 0 & p & 0 \\ 0 & m & 0 & -p \\ p & 0 & -m & 0 \\ 0 & -p & 0 & -m \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{pmatrix} = E \begin{pmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{pmatrix}.$$

## 19. Klein Tunneling in graphene

1+3+6 Points




Figure 1: Left: Schematic drawing of a Dirac electron incident on a potential barrier. Right: Definition of the angles used in the problem. Assume that the sample is infinite in the  $y$ -direction.

Consider a Dirac electron with energy  $E$  incident on a potential barrier of size  $V$  as shown in the figure.

- Why is it sufficient to only require continuity of the wave-function and not its derivative?
- Assume the electron is incident at some angle  $\phi$  in regions I and III and  $\theta$  in region II, such that  $k_x = k \cos \phi$ ,  $k_y = k \sin \phi$  in regions I and III, while  $\theta = \arctan(k_y/q_x)$  with  $q_x = \sqrt{(V - E)^2/v^2 - k_y^2}$  and  $v = |\mathbf{k}|/m$  in region II. Explain why the wave-functions in the different regions can be written as

$$\begin{aligned}\psi_{\text{I}}(x) &= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ s e^{i\phi} \end{pmatrix} e^{i(k_x x + k_y y)} + \frac{r}{\sqrt{2}} \begin{pmatrix} 1 \\ s e^{i(\pi - \phi)} \end{pmatrix} e^{i(k_y y - k_x x)}, \\ \psi_{\text{II}}(x) &= \frac{a}{\sqrt{2}} \begin{pmatrix} 1 \\ s' e^{i\theta} \end{pmatrix} e^{i(q_x x + k_y y)} + \frac{b}{\sqrt{2}} \begin{pmatrix} 1 \\ s' e^{i(\pi - \theta)} \end{pmatrix} e^{i(k_y y - q_x x)}, \\ \psi_{\text{III}}(x) &= \frac{t}{\sqrt{2}} \begin{pmatrix} 1 \\ s e^{i\phi} \end{pmatrix} e^{i(k_x x + k_y y)}.\end{aligned}$$

Here  $s = \text{sgn}(E)$  and  $s' = \text{sgn}(E - V)$ . What is the physical significance of  $r$ ,  $a$ ,  $b$ , and  $t$ ?

- Use the continuity of the wave-function to calculate the transmission through the barrier  $T(\theta, \phi, Dq_x) = |t|^2$ . What do you get for  $Dq_x = n\pi$  with  $n$  integer? For general values of  $Dq_x$ , investigate what happens when  $\phi, \theta \rightarrow 0$ .

*Hint: You might want to use a computer algebra system to solve the resulting linear equation system for  $t$ , and to compute  $|t|^2$ .*