

## Advanced Statistical Physics - Problem Set 5

*Summer Term 2020*

**Due Date:** Tuesday, May 12, 10:00 a.m., solutions must be mailed to [stp2.leipziguni@gmail.com](mailto:stp2.leipziguni@gmail.com)

### 1. Coupling to a massless field

*2+2+2+2+2+2+2 Points*

Consider an  $n$ -component vector field  $\vec{m}(x)$  coupled to a scalar field  $A(x)$ , through the effective Hamiltonian

$$\beta\mathcal{H} = \int d^d x \left[ \frac{K}{2} (\nabla \vec{m})^2 + \frac{t}{2} \vec{m}^2 + u(\vec{m}^2)^2 + e^2 \vec{m}^2 A^2 + \frac{L}{2} (\nabla A)^2 \right],$$

with  $K$ ,  $L$ , and  $u$  positive.

a) Assume  $\vec{m}(x) = \bar{m} \hat{e}_\ell$  and  $A(x) = \bar{A}$ , and find the saddle point solution  $\bar{m}$  for  $t > 0$  and  $t < 0$ .

b) Sketch the heat capacity  $C = \partial^2 \ln Z / \partial t^2$  in the saddle point approximation, and discuss its singularity as  $t \rightarrow 0$

c) Include fluctuations by setting

$$\begin{cases} \vec{m}(x) = (\bar{m} + \phi_\ell(x)) \hat{e}_\ell + \phi_i(x) \hat{e}_i, \\ A(x) = a(x), \end{cases}$$

and expanding  $\beta\mathcal{H}$  to quadratic order in  $\phi$  and  $a$ . Hint: after substituting the above in  $\beta\mathcal{H}$ , the linear terms vanish at the minimum and the second order terms give

$$\begin{aligned} \beta\mathcal{H}_2 = & \int d^d x \left[ \frac{K}{2} (\nabla \phi_\ell)^2 + \frac{t + 12u\bar{m}^2}{2} \phi_\ell^2 \right] + \int d^d x \left[ \frac{K}{2} (\nabla \phi_t)^2 + \frac{t + 4u\bar{m}^2}{2} \phi_t^2 \right] \\ & + \int d^d x \left[ \frac{L}{2} (\nabla a)^2 + \frac{2e^2\bar{m}^2}{2} a^2 \right] + \mathcal{O}(\phi^3). \end{aligned}$$

d) Use your results from (c) to find the correlation lengths  $\xi_\ell$ , and  $\xi_t$ , for the longitudinal and transverse components of  $\phi$ , for  $t > 0$  and  $t < 0$ .

e) Find the correlation length  $\xi_a$  for the fluctuations of the scalar field  $a$ , for  $t > 0$  and  $t < 0$ .

f) Compute the correction to the saddle point free energy  $-\ln Z/V$ , from fluctuations. (You can leave the answer in the form of integrals involving  $\xi_\ell$ ,  $\xi_t$ , and  $\xi_a$ ). Hint: Perform a Fourier transform in the Hamiltonian beta H2 to obtain Gaussian integrals over  $D[\phi_t]$ ,  $D[\phi_\ell]$ , and  $D[a]$ . Then compute the Gaussian integrals to obtain an expression for  $Z$ . You do not need to perform the sums over momenta that you obtain after the Fourier transformation.

g) Find the fluctuation corrections to the heat capacity in (b), again leaving the answer in the form of integrals.