

# 1 Relaxation time estimate for Lévy flights in a confined two dimensional region

One can estimate the time to reach equilibrium for a two-dimensional Lévy flight in a confined region of linear extent  $L$  as follows. Assume that the single step radial distribution of the random walk can be approximated by

$$p_{\Delta t}(\mathbf{x}) = (1 - Q)\delta(\mathbf{x}) + Q f_{\delta L}(\mathbf{x}). \quad (1.1)$$

Here  $\Delta t$  denotes the typical time between single steps,  $Q$  the fraction of walkers which jump a distance  $d > \delta L$  and  $(1 - Q)$  the fraction which remains in a disk defined by  $|\mathbf{x}| \leq \delta L$ . The function  $f_{\delta L}(\mathbf{x})$  comprises the power-law in the single steps, characteristic for Lévy flights:

$$f_{\delta L}(\mathbf{x}) = C \delta L^\beta |\mathbf{x}|^{-(1+\beta)} \quad |\mathbf{x}| \geq \delta L.$$

Inserting this into Eq. (1.1) one obtains that  $f_{\delta L}(\mathbf{x})$  is normalized to unity and that the normalization constant  $C$  is independent of the microscopic length  $\delta L$ . The Fourier-transform of  $p(\mathbf{x})$  is given by

$$\tilde{p}(\mathbf{k}) = (1 - Q) + Q \tilde{f}_{\delta L}(\mathbf{k}).$$

The Fourier-transform of the probability density function  $W_N(\mathbf{x})$  of the walker being located at a position  $\mathbf{x}$  after  $N$  steps can be computed in terms of  $\tilde{p}(\mathbf{k})$  according to

$$\tilde{W}_N(\mathbf{k}) = \tilde{p}(\mathbf{k})^N \approx \left(1 - Q \delta L^\beta |\mathbf{k}|^\beta\right)^N \approx e^{-Q N |\delta L \mathbf{k}|^\beta}. \quad (1.2)$$

The relaxation time in a confined region is provided by the lowest mode

$$k_{\min} = \frac{L}{2\pi}.$$

Inserted into (1.2) with  $N = t/\Delta t$  one obtains

$$T_{\text{eq}} \approx \delta T / Q (L/2\pi\delta L)^\beta.$$